



# Modelling responses of forages to climate change with a focus on nutritive value

P. Virkajärvi, P. Korhonen, G. Bellocchi, Y. Curnel, L. Wu, G. Jégo, T. Persson, M. Höglind, M. Van Oijen, A.-M. Gustavsson and R.P. Kipling





# Nutritive value of forage is key factor affecting

- ruminant nutrition
- animal performance
- need for other feeds
- productivity of the system
- quality of final products
- environmental impact of production

# Climate change is expected to affect the nutritive value of grasslands by

- affecting plant physiological processes
- via effects on species composition.

Kipling et al 2016. Modeling European ruminant production systems: Facing the challenges of climate change. *Agricultural Systems* 147: 24-37





# Variability in the grassland based systems



© L Frondelius

Frequency of defoliation, nutrient cycling



© P Virkajärvi



© P Virkajärvi

Monoculture



© P Virkajärvi

Binary mixture



Multispecies



© P Virkajärvi



# Variables used to describe the nutritive value (NV) of forage grass based feeds

## Energy variables

- ME:** metabolizable energy
- NEL:** net energy of lactation
- NEM:** net energy of maintenance
- FME:** fermentable metabolisable energy
- FEm:** feed unit for milk production (Norwegian)
- ...

## Digestibility variables

- CWD:** cell wall digestibility
- CWC:** cell wall content/concentration
- IVCWD:** in vitro cell wall digestibility
- NDF:** neutral detergent fiber
- NDS:** neutral detergent solubles
- dNDF:** in vitro digestibility of NDF
- iNDF:** indigestible NDF
- pdNDF:** potentially digestible NDF
- OMD:** organic matter digestibility
- DOM:** digestible organic matter
- IVOMD:** in vitro organic matter digestibility
- IVTD:** in vitro true digestibility of dry matter
- TDN:** total digestible nutrients
- D-value:** concentration of digestible organic matter in DM
- ...

## Protein variables

- N concentration**
- CP:** crude protein
- DCP:** digestible crude protein
- RDP:** rumen digestible protein
- ERDP:** effective rumen degradable protein
- ADIP:** acid detergent insoluble protein
- DUP:** digestible undegradable protein content
- ADIN:** acid-detergent insoluble nitrogen
- ...

McDonald et al 2002. Animal Nutrition.

- Dissimilar feeding systems


- Most important are:
  - energy value
  - protein content





# Factors and processes behind NV to be modelled

- Botanical composition
  - Grass - legumes- dicots
    - Grass functional traits
- Tiller dynamics & phenological stage
  - senescence
- Leaf to stem ratio
- Proportion of cell wall
- Chemical composition
  - Lignification of cell walls
  - in general (N, minerals etc.)





## The aim

- To review the extent to which current process-based grass growth models are capable of characterising the nutritive value of forage grasses in relation to the projected climate change.
- To identify of modelling approaches, the key characteristics of the forages and the production systems the individual models are developed for





# The models

- Including process based growth models (PBMs) of temperate climate
- Reviewed from literature
- A questionnaire survey was sent to MACSUR knowledge hub partners





# Results

- 8 PBMs simulating forage NV were included in the study
  - 6 from Europe
  - 2 from USA and Canada
- Developed mainly for silage - 2 models includes grazing
- Both generic and species specific models





# The models

Acronym	Name	Developer/owner	First version (year)
<b>BASGRA</b>	BASic GRAss model	Centre for Ecology and Hydrology, UK; Norwegian Institute of Bioeconomy Research , Norway	2016
<b>CATIMO</b>	Canadian Timothy Model	Natural Resources Institute Finland (Luke)	2002
<b>IFSM</b>	Integrated Farm System Model	US Department of Agriculture, Agricultural Research Service (USDA ARS, USA)	2005 (first reference)
<b>MCPy</b>	Modélisation de la Croissance des prairies	Centre Wallon de Recherches agronomiques	2001
<b>PaSim</b>	Pasture Simulation model	French National Institute for Agricultural Research (INRA)	1998 (full description)
<b>Qual</b>	Integrated Dynamic Model	Swedish University of Agricultural Sciences	1994
<b>SPACSYS</b>	Soil Plant and Atmosphere Continuum SYStem	Scottish Agricultural College and Rothamsted Research, UK	2007 (first reference)
<b>STICS</b>	Simulateur mulTidisciplinaire pour les Cultures Standards	Institut National de la Recherche Agronomique (INRA, France)	1998



# The main variables used to describe the NV of forages

Model	Reference	Monoculture/ Mixture	Energy variables		Digestibility variables		CP variables	
			ME	other	NDF	other	[N]	other
BASGRA	(Höglind <i>et al.</i> 2016)	Monoculture	-	-	-	-	Yes	CP
CATIMO	(Bonesmo and Bélanger, 2002)	Monoculture	-	-	Yes	IVTD, dNDF	Yes	RNC
IFSM	(Rotz <i>et al.</i> , 2015)	Mixture	Yes	NEL/ NEM	Yes	TDN	Yes	CP
MCPy	(Stilmant <i>et al.</i> , 2001)	Mixture	-	VEM	Yes	-	-	-
PaSim	(Graux <i>et al.</i> , 2011)	Mixture	MEI	NELh	Yes	OMD, dNDF	Yes	-
QUAL	(Gustavsson <i>et al.</i> , 1995)	Monoculture	Yes	-	-	OMD, IVTD	Yes	CP
SPACSYS	(Wu <i>et al.</i> , 2007)	Mixture	-	-	-	-	Yes	-
STICS	(Brisson <i>et al.</i> , 1998; Jégo <i>et al.</i> , 2013) for NV	Mixture	-	-	Yes	IVTD, dNDF	Yes	CP



CP, crude protein; dNDF, in vitro digestibility of NDF; IVTD, in vitro true digestibility of dry matter; ME, metabolizable energy; MEI, metabolizable energy intake; [N], nitrogen concentration of forage; NDF, neutral detergent fiber; NEL, net energy of lactation; NELh, net energy content of the ingested herbage; NEM, net energy of maintenance; OMD, organic matter digestibility; RNC, relative nitrogen concentration; TDN, total digestible nutrients, VEM, available energy for milk production.



# Discussion (1)

- PBMs tend to be fairly comprehensive in their consideration of weather variables that are expected to change
  - Limitation in CO<sub>2</sub> effect on water use efficiency, sugars, N, etc.
- Variables describing NV are clearly the most essential ones
  - DM of OM digestibility,
  - NDF, NDF digestibility
  - CP,
- Variables are very useful for feed planning.





## Discussion (2) - improvements


### How to simulate:

1. physiological adaptation of the plants to changes in environmental conditions
2. formation and senescence of tillers
3. the dynamics of leaf chemical composition including water soluble carbohydrates
4. the response of C and N allocation to environmental change
5. quantify the relative importance of grazing regime and harvest dates
6. use information from field and laboratory trials with different genotypes to parameterize alternative cultivars, e.g. to represent developmental stages
7. improve the link between plant and soil water and soil N



# Conclusions


- This work is the first step towards gathering and clarifying information about the possibilities of modelling NV
- The number of PBMs capable to predict NV is rather limited
- PBMs tend to be fairly comprehensive in their consideration of weather variables - reaction and acclimation to CO<sub>2</sub> level rise
- Development needed to reduce uncertainties
- a strong need for data including frequent time series of forage NV from experiments mimicking global change conditions
  - sharing of existing data sets



# THANKS!



Modelling European Agriculture with Climate Change for Food Security  
— a FACCE JPI knowledge hub —



FACCE JPI

MACSUR

[www.macsur.eu](http://www.macsur.eu)