

FACCE-MACSUR

D-L3.3. Synergies between mitigation and adaptation to Climate Change in grassland-based farming systems

Del Prado A¹., Van den Pol-van Dasselaar, A²., Chadwick, D³., Misselbrook, T⁴., Sandars, D⁵., Audsley, E⁵. and Mosquera-Losada, M.R⁶

¹Basque Centre for Climate Change (BC3), Alameda Urquijo, 4, 4^o-1^a /48008 Bilbao, Spain

²Wageningen UR Livestock Research, 8219PH Lelystad, The Netherlands

³School of Environment, Natural Resources and Geography, Bangor University, Bangor, Wales, UK

⁴Rothamsted Research, North Wyke, Okehampton, Devon, EX20 2SB, UK

⁵School of Applied Sciences, Cranfield University, Cranfield, Bedford MK43 0AL

⁶Departamento de Producción Vegetal, Universidad de Santiago de Compostela, Spain

*agustin.delprado@bc3research.org

Instrument:	Joint Programming Initiative
Topic:	Agriculture, Food Security, and Climate Change
Project:	Modelling European Agriculture with Climate Change for Food Security (FACCE-MACSUR)
Due date of deliverable:	month 32
Start date of project:	1 June 2012
Duration:	36 months
Theme, Work Package:	LiveM 1
Deliverable lead partner:	BC3
Deliverable reference num.:	D-L3.3
Deliverable type:	Report/Workshop/Action
Confidential till:	Not applicable

Revision	Changes	Date
1.0	First Release	2015-01-23

Abstract/Executive summary

Climate change mitigation and adaptation have generally been considered in separate settings for both scientific and policy viewpoints. Recently, it has been stressed (e.g. by the latest IPCC reports) the importance to consider both mitigation and adaptation from land management together. To date, although there is already large amount of studies considering climate mitigation and adaptation in relation to grassland-based systems, there are no studies that analyse the potential synergies and tradeoffs for the main climate change mitigation and adaptation measures within the current European Policy context. This paper reviews which mitigation and adaptation measures interact with each other and how, and it explores the potential limitations and strengths of the different policy instruments that may have an effect in European grassland-based livestock systems.

Table of Contents

1. Introduction	2
2. Climate Change mitigation now and in the future in grassland-based livestock systems	2
3. Main climate change effects on European grassland-based livestock systems	3
4. Potential synergies and tradeoffs between strategies to adapt to Climate Change and measures to mitigate GHG emissions	4
4.1. Climate change affecting feed supply (grazing and forage)	4
4.2. Climate change affecting feed supply (purchased feed and different crop rotations).....	5
4.3. Climate change affecting feed supply (use of by-products and alternative forages)	6
4.4. Changes in fertiliser management, diet and genetics to increase N use efficiency	7
4.5. Soil management, plant biodiversity and new plant breeds to improve system resilience against environmental stress conditions and prevent soil erosion	7
4.6. Agroforestry systems	8
5. Policy implications	9
6. References	12

1. Introduction

In the last IPCC report (AR5-WGIII: IPCC, 2014b), for the first time, most of the terrestrial land comprising agriculture, forestry and other land use (AFOLU) was considered altogether. Moreover, it was also highlighted in the AFOLU chapter (IPCC, 2014b) the importance to consider the systemic feedbacks and interactions between mitigation and adaptation options from land management (Separate sub-section: 11.5). In grassland-based systems, however, the potential interactions between mitigation and adaptation options, compared with forest or arable systems, have received much less attention and this has been reduced to changes in carbon (C) stocks and pasture productivity. Changes in biogeochemical cycles (mainly C and N) and water cycles are expected to exert large impacts on livestock productivity and N and C emissions from grassland-based systems (i.e. CO₂, CH₄ and N₂O). Climate change impacts on livestock will include effects of forage and feed quality and productivity, direct impacts of changes in temperature and water availability on animals, and indirectly through livestock disease increase (IPCC, 2014a). However, socio-economic changes are expected to have a still greater effect on mitigation and adaptation potentials (Schmidhuber and Tubiello, 2007).

Climate mitigation options in grassland systems mainly include practices that increase soil C stocks or can help to reduce GHG (greenhouse gases) emissions at the soil, feed, animal or manure management level. However, at a wider context, demand-side measures (e.g. human dietary changes, reducing losses and wastes in the agro-food chain) or substitution of fossil fuels by biomass can also play an important role in mitigating climate change. Mitigation options in grassland-based systems need also to be addressed for their potential impact on all other ecosystem/environmental services provided by grasslands for current and future scenarios, as climate regulation is just one of the services amongst a varied list (e.g. food production). Mitigation and adaptation in grassland-based systems are closely integrated through a network of feedbacks, synergies and risk of trade-offs. Mitigation measures may also be vulnerable to climate change or there may be possible synergies and trade-offs between mitigation and adaptation options (IPCC, 2014b).

Many policies are directly (e.g. Kyoto Protocol) and indirectly (e.g. Common Agricultural Policy: CAP) affecting the potential success of implementing measures that both reduce GHG emissions and help to adapt grassland-based livestock systems to Climate Change. There is however a need for a more integrated and scientifically-based regulatory approach.

Although there are already many studies that have reviewed measures regarding grassland-based systems on GHG mitigation (e.g. Smith et al., 2007; Verge et al., 2007) or Climate Change adaptation (e.g. Bryan et al., 2009; Olesen et al., 2011; Tingem et al., 2009), only few studies have assessed the benefits and trade-offs of their synergistic effects (e.g. Lal et al., 2011).

The main objective of this paper is to provide a high-level assessment of synergies and trade-offs for the main potential Climate Change mitigation and adaptation measures in grassland-based livestock systems within the current European Policy context.

2. Climate Change mitigation now and in the future in grassland-based livestock systems

The main aim of the mitigation options in grassland-based systems is to reduce emissions of CH₄ or N₂O and/or to increase soil C storage, especially by soil as grasslands account for 75% of C in the terrestrial ecosystems (Lal, 2005; Dresner et al., 2007). Recently, more or less comprehensive reviews on GHG mitigation from grassland-based systems have been produced (e.g. Project ANIMALCHANGE: Van den Pol-van Dasselaar, 2012; UNEP, 2013;

Havlík et al., 2014; Del Prado et al., 2013a). Nitrous oxide is formed in the soil through nitrification and denitrification (Wrage et al., 2001) and controlled by a number of site-specific factors, including soil moisture content (Del Prado et al., 2006), temperature (Dobbie et al., 2001) and also, management factors such as fertilizer (Cardenas et al., 2010) and management of soil organic matter content (Mosquera-Losada et al., 2011a) and grazing (Van den Pol-van Dasselaar et al., 2008). Carbon sequestration also depends on edaphoclimatic conditions (Theng et al., 1989), the presence of trees (Mosquera-Losada et al., 2011a) and the organic matter quality and quantity (Mosquera-Losada et al., 2011a and 2011b). Methane can be produced via enteric fermentation, which depends greatly on the level of feed intake, the quantity of energy consumed and feed composition, or can be produced at the manure management level, which increases with temperature, and with increased biodegradability of the manure (Monteny et al., 2006).

As highlighted in the last IPCC report on mitigation of Climate Change in the AFOLU sector, there is an emerging scientific activity on prediction of the likely impact of the Climate Change on the potential to reduce net GHG emissions (i.e. impacts on N_2O and CH_4 emissions and on the rates of C sequestration) in the AFOLU sector in general, and in grassland-based systems in particular. Mitigation options available today in the grassland-based farming sector may not be available or as effective with further global warming. Soil C storage has been shown to be vulnerable not only to climate change but more importantly, to changes in the disturbance regime, both natural and human-induced. Land use projections indicate large changes in land use and potentially this change will exacerbate the release of CO_2 from soils including grasslands. Increasing temperatures, when water is not limiting, are expected to accelerate soil organic matter (SOM) decomposition rates but result in an increase of C returns through plant residues considering the CO_2 fertiliser effect and the lengthening of the growing season (Bindi and Olesen, 2011). Increasing SOM will enhance soil C storage and may also increase above and belowground biomass production or at least improve yield stability (Pan et al., 2009). However, biological processes resulting in N_2O emissions (i.e. denitrification) could be stimulated by greater SOM. Moreover, increased variability and higher frequency of extreme events will negatively impact soil C storage, by both decreasing production levels and enhancing soil C losses. At the manure level, GHG emission changes are expected in relation to temperatures and sometimes indirect effects driven by changes in the composition of the feed (e.g. digestibility).

3. Main climate change effects on European grassland-based livestock systems

During the last century, the climate in Europe has changed more than in other areas of the world (IPCC, 2007). Compared to the pre-industrial era, when the mean annual temperature increased by 0.8°C globally, it increased by 1.2°C in Europe. Based on theoretical models, a further increase of $1.0\text{--}5.5^\circ\text{C}$ is expected by the end of the twenty-first century (Christensen et al., 2007). The increase in temperature has been most apparent in mountainous areas such as the Alps, which tend to have high biodiversity and where temperature increased by 2°C during the twentieth century (EEA, 2009). This is twice the average temperature increase for the northern hemisphere. In addition, the quantity and distribution of precipitation have also changed in Europe during the twentieth century. Although there has been a 20% decrease in rainfall in southern Europe, there has been a 10-40% increase in rainfall in northern Europe. Furthermore, an increase in the frequency of extreme weather events is predicted across the European continent (EEA, 2008).

The most important impacts of climate change on grassland-based farming systems in Europe are expected to be through changes in pasture productivity and forage quality,

therefore potentially affecting the duration of the grass growing season and the forage supply to ruminants. An example about how different intra-year temperature and precipitation regimes affect total and also seasonal distribution of pasture was found by Mosquera-Losada and González-Rodríguez (1998) in dairy systems. This paper highlights the importance of having flexible grazing systems, which affects annual and instantaneous stocking rate. The intensification of the hydrological cycle caused by more intensive rainfall and longer dry periods is expected to result in higher risks of soil erosion and nutrient leaching in currently wet temperate climates. Changes in precipitation patterns in drier areas will lead to higher dependency on stored soil moisture storage and seasonality for supporting grass growth. Elevated concentrations of CO₂ may also increase water use efficiency through reduced plant stomata aperture, but increase run-off risk through reduced plant transpiration, thus resulting in excess water at the land surface (Betts et al., 2007). Biological and physical processes regulating nutrient cycling in grassland-based farming systems may actually be more sensitive to extreme events rather than changes in average climatic conditions. For Europe, an increased risk of low forage production in summer due to severe summer droughts events is expected to be offset by the appearance of new opportunities for forage production in other seasons due to warming effects. For southern latitudes, higher evapotranspiration rates will negatively affect grass yield and the period of grass growth will be shorter unless the grassland is irrigated. In general, poorer grass nutritional qualities, e.g. lower grass digestibility, can also be expected.

4. Potential synergies and tradeoffs between strategies to adapt to Climate Change and measures to mitigate GHG emissions

GHG emission mitigation choices may further enhance or reduce resilience to climate variability and change in terms of ecosystem goods and services provision, and thus influence the potential of grassland-based systems to adapt to Climate Change. Climate change may affect climate adaptation and mitigation strategies through changes in feed supply, animal diet composition, animal and plant breeding, soil management, enhancement of floral biodiversity and via more resistant and resilient production systems against climate change (e.g. agroforestry systems).

4.1. Climate change affecting feed supply (grazing and forage)

Spring growth, provided that water resources for grass growth are available, and winter production may benefit from mild climate conditions. This can contribute to improve the farm's degree of forage autonomy and security of livestock systems when facing more hazardous climate conditions (e.g. summer droughts) through the extension of the grazing season and the reduction of forage requirements (Graux et al., 2013). For example, forage resource usually stored for over-wintering livestock could be partially redistributed in summer to deal with increased risk of forage deficits (Graux et al., 2013). However, for southern latitudes and dates getting closer to the XXI century (e.g. UK: Del Prado et al., 2009) the projections suggest that grazing activity will be constrained due to too high temperatures and excessive drought in Europe. Extending grazing seasons by e.g. the presence of shelter/shade belts of trees would reduce the wind speed and therefore evapotranspiration (ETP). The presence of trees at low density would also increase the duration of the growing season due to their presence, which may partly reduce GHG emissions (Tackas and Frank, 2009) through improving soil N recovery by trees, but may also become hot-spots for N₂O from overlapping urine patches, and soils could become eroded due to the action of hooves in camping areas used by livestock for shelter/shade. Extending the grazing season in some cases may also be limited by the bearing capacity of the soil driven by good soil structure degradation (e.g. poaching caused by trampling cattle or/and severe summer droughts, etc.) and therefore, it may, in some cases be impractical. Hence, avoiding compaction by traffic, tillage (Pinto et al., 2004) and grazing livestock (De

Klein and Ledgard, 2005) may help to maintain grasslands in good conditions and also to reduce N₂O emissions.

Poorer grass nutritional qualities, e.g. lower grass digestibility, will lead to higher CH₄ emissions from enteric fermentation of cattle (Hart et al., 2009). Although if lower forage quality would mean reduced livestock ‘yields’ and/or quality then market reaction would be to import feeds. For those systems extending the grazing season (e.g. Figure 1: UK: Misselbrook et al., 2013), smaller volumes of manure storage could lead to a reduction in CH₄ emissions from manure handling. However, the rise in average and extreme temperatures, should the frequency of manure removal from the storage remain unchanged or no additional structural measures are implemented to either lower the manure storage temperature or to aerate the slurry, could increase the amount of CH₄ per kg of volatile solid of manure and therefore, lead to similar total CH₄ emissions from manure management. Moreover, even if the CH₄ emissions from manure management were smaller, this value (in CO₂ equivalents) would be partially offset or accentuated by an increase in N₂O emissions promoted by the increase in grazing activity.




Figure 1. Comparison between adapted (extending one month grazing) and un-adapted typical dairy farms in the south west England (2020) for GHG, NO₃- leaching and NH₃ emissions. Values for the adapted scenario <1 indicate a reduction in emissions (adapted from Misselbrook et al., 2013).

4.2. Climate change affecting feed supply (purchased feed and different crop rotations)

Changes in grassland productivity will affect either animal productivity or the amount of purchased feed required (Mosquera-Losada and González-Rodríguez, 1998). For semi-arid regions (e.g. south of Europe), a reduction in annual grass productivity will lead to lower animal productivity or will have to be compensated with a larger share of imported feedstock with associated monetary and environmental costs, which may translate into a potential loss of resilience in grassland-based livestock systems. In some of these regions tree presence to feed animals with acorns, could supply part of these needs (Moreno and Pulido, 2009).

An increase in the establishment of rotations best suited to the area or crop rotations with legumes annual crops (Bryan et al., 2011) may also occur as an adaptation strategy. Some crops that currently grow mostly in southern Europe will become more suitable further north or in higher altitudes areas in the South. For example, forage maize, may become more common across in the boreal regions of Europe. Maize forage, however, tends to make the management system less flexible to inter-annual temperature/precipitation variations.

Moreover, maize area cannot be used for grazing during the summer or autumn if no grass is available during this period. In contrast, grass areas can be open or harvested for silage if a restriction or an excess of grass production happens (Mosquera-Losada and González-Rodríguez, 1998). At the animal level, forage maize animal intake is generally promoted at the expense of grass due to a better balance between protein intake and soluble carbohydrates (e.g. through increasing starch concentration in the diet), which additionally may help to increase animal energy use efficiency and decrease CH₄ emissions per kg DM intake. However, this CH₄ reduction may be offset by larger N₂O emission losses and a larger CO₂ release of converting some grassland into arable land (Vellinga and Hoving, 2011).

Conversely, converting crops to pasture has been found to reduce N₂O emissions (Eagle et al., 2012) and also contribute to sequester soil C, especially in the first years after conversion. Leguminous species are well adapted to future conditions of climate change (Kreyling et al., 2012) considering that their optimum temperature is higher than non-leguminous crops and that they also have more positive responses to elevated concentrations of CO₂ (Soussana and Lüscher, 2007) than non-legume species. In a situation with a larger share of mixed legume/grass pastures, in addition to presenting climate adaptive advantages over conventional pastures, these systems have lower requirements for N fertilizer through the use of biological N fixation of nodules on the roots of legumes, which would lead to energy savings and GHG emissions reductions from both fertilizer production and use (Del Prado et al., 2011; Zhang et al., 2013).

4.3. Climate change affecting feed supply (use of by-products and alternative forages)

Different by-products from agricultural, forestry, agro-industry and bioenergy activities can also be used for feeding ruminants as an adaptive response to forage supply seasonal constraints. Rinne et al. (2012) reviewed different by-products (e.g. camelina meal, tomato pomace) that are currently underutilized but that could potentially be used as feed for low input and organic dairy production systems. Those practices are currently used as part of some livestock systems at a regional level (Correal et al., 2009). These by-products vary in their geographical availability, nutritional value, their effect on rumen CH₄ and N excretion (i.e. effect on GHG mitigation) and have logistic-related challenges.

Environmentally speaking (e.g. GHG intensity), the use of some of these by-products as animal feed may not always be the best option in comparison with their use in bioenergy or for soil improvement. In this sense, removal of crop residues from cropping systems for use in bioenergy, if this means that soil C contents are being depleted (e.g. straw: Liu et al., 2014), will bring large risks of negative impacts on adaptation measures and potentially, small or negligible positive effects on the reduction of net GHG emissions. Mitigation and adaptation conflicts may therefore appear as one chooses a particular use of the by-product or another.

Other alternative forage supply may include tree leaves and shrubs, particularly in small-scale livestock farms with dry to semi-arid climates. Such species can alleviate feed shortages, or even fill feed gaps in the winter and especially in the summer, when grassland growth is limited or dormant due to unfavourable weather conditions

(Papanastasis et al., 2008). Although some species have leaves with a low content of CP and a high content of fiber and contain high levels of secondary compounds such as tannins, alkaloids, saponins and oxalates which reduce the nutritive value of poor-quality diets, some of these compounds (e.g. condensed tannins), when improved temperate forages are fed, can also have substantial benefits for ruminant productivity (i.e. reducing CH₄) and health (Waghorn and McNabb, 2003). Moreover, there are also other species (e.g. *Morus alba*, *Fraxinus excelsior*, *Betula alba*) whose young leaves are rich sources of protein and fibre and generally used in the past to feed animals before modern techniques like fertilizer were used.

4.4. Changes in fertiliser management, diet and genetics to increase N use efficiency

Manipulating the diet (e.g. feeding nitrification inhibitors: Ledgard et al., 2008 or salt supplementation) during the grazing period has also been proposed as a means to reduce N₂O emissions. Improving fertiliser efficiency, optimising methods, timing and rates of applications (Brown et al., 2005), using NH₄⁺-based fertilisers rather than nitrate-based ones (e.g. Dobbie and Smith, 2003) and employing nitrification chemical inhibitors (e.g. Zaman et al., 2009) may also have a role in both mitigation (i.e. reduction of direct and indirect soil N₂O emissions) and adaptation (through a better N use efficiency at the soil-plant level).

New traits in animals and grasses may also assist farmers to both mitigate and adapt to Climate Change. Del Prado and Scholefield (2008), for example, using a farm modelling approach, evaluated the scope for different animal and plant genetic traits, some existing and other theoretical, to help reduce GHG emissions on UK dairy farms. More efficient animals in utilising N (Alford et al., 2006) have also been proposed to decrease the impact of urinary N during grazing. Some of the traits, e.g. improved N use efficiency in grasses (e.g. high sugar grasses: Wilkins et al., 2000) could actually be both potentially useful for Climate mitigation and may also promote Climate adaptation as they may reduce GHG emissions from urine-related N₂O emissions and improve the quality of the forage, which may be beneficial in future scenarios where climate has a detrimental effect on grass nutritional properties.

4.5. Soil management, plant biodiversity and new plant breeds to improve system resilience against environmental stress conditions and prevent soil erosion

Other strategies to both mitigate and adapt to Climate Change may involve management practices that target directly to the soil, both improving the capacity to store water and to prevent soil erosion. By increasing the ability of soils to hold soil moisture and to better withstand erosion by enriching biodiversity through more diversified cropping systems, grassland systems will be able to sequester more soil C and also to better resist extreme events such as droughts and /or floods, both of which are projected to increase in frequency and severity in future warming climates (Rosenzweig and Tubiello, 2007). The measures for the conservation of soil moisture may also include changes in tillage practices. Reduced tillage, for example, increases the resilience to climate change through improved soil fertility and increased capacity for water retention in the soil. This improvement is expected in the long-term productivity potential when tillage is reduced (Olesen et al., 2011). The reduced tillage at pasture reseeding promotes C sequestration and preservation in pastures and is considered to be more effective under conditions of water deficit (Alvaro -Fuentes et al., 2011). It leads also to significant savings in CO₂ emissions produced by machinery. However, the impact on N₂O emissions under different conditions is unclear (Estavillo et al, 2002; Pinto et al, 2004). Nitrous oxide emissions appear to be strongly influenced by soil water content immediately after nitrogen

fertilization (Del Prado et al., 2006). In view of this dominant effect of a particular soil moisture level coinciding with tillage and fertilization, it is key to find the best timing for the renewal of pasture. Velthof et al. (2010), for example, considering average Dutch climatic season conditions, suggest that this pasture renewal should take place in spring rather than fall because Dutch autumn, compared with spring is generally wetter and N uptake by the reseeded grass is lower. The effect of reduced tillage has also been observed by increasing the periods between which a pasture is renewed. Vellinga et al. (2004), for example, found that although tillage increased N₂O and CO₂ in the intensively managed pastures in the studied year, in the long run, the renovation of the pastures was more important to prevent the deterioration in pasture quality and thus, to prevent from soil loss and large productivity losses.

For areas which are subject to severe or extremely severe environmental stress conditions the establishment of a community of pastures formed by species that ensure ecological stability, both in ecosystem resistance and resilience, is key as an adaptation measure to Climate Change (Volaire et al., 2014). Additionally the species composition of the pasture is expected to undergo changes, as for example, warming will favour C4 species over C3 species (Howden et al., 2008). Biodiversity should act as a safeguard of ecosystem functioning, thus promoting a more stable ecosystem to avoid fluctuations arising from adverse climatic fluctuations (Volaire et al., 2014). Promoting biodiversity could also have an effect on the mitigation potential of pastures and in some occasions of rumen methane. Considering that N remains one of the main elements that determines the diversity of plants, the application of less fertilizer should be a requirement to increase diversity in different floral species in grasslands (Mountford et al., 1993). This reduced input fertilizer would be necessarily associated with lower emissions of N₂O per ha and potentially a greater amount of C accumulated in the soil.

New grass breeds have already been tested to improve water use efficiency. For example, McLeod et al. (2013) tested in the UK a novel grass *Festulolium* hybrid capable to reduce runoff by 40-50% compared to a leading UK nationally recommended *L. perenne* cultivar and *F. pratensis* over a two year field experiment. The rapid growth and turnover of roots in the hybrids resulted in greater soil water storage capacity in the plots with observed lower rainfall runoff. This may, in turn, have significant effects on N₂O emissions and soil C storage.

4.6. Agroforestry systems

Agroforestry is a well-founded example of mitigation and adaptation synergy (e.g. IPCC, 2014b; EU forest strategy: EU, 2013) since trees planted and grassland soils sequester C and tree and grassland products provide livelihood to communities, especially during drought years (Verchot et al., 2007). Agroforestry in general and silvopastoral systems in particular lead to greater resilience to climate change due to improved soil conditions and management efficiency in water use (Kumar et al., 2011). Its characteristics are able to reduce evapotranspiration and thus improve the maintainability of soil water (Tackas and Frank, 2009). These practices also have a great potential to offset GHG emissions through the sequestration of C in soil and tree biomass and avoiding the release of NO₃ leaching (indirect N₂O emissions) (Rigueiro et al., 2009). Moreover, these systems also improve the N use efficiency of the system and offer large resilience against climate change stress conditions through the reduction of temperature of the system (Rigueiro et al. 2009). It can also help reduce erosion of adjacent fields handled more intensely (Verge et al., 2007).

5. Policy implications

Climate mitigation policies and measures may exhibit synergies and risk trade-offs with climate adaptation (Bates et al., 2008). However, policies of mitigation and adaptation are often being considered in separate settings, resulting in potential conflicts. An integrated adaptation and mitigation framework is important to ensure that trade-offs between the two are minimized and synergies encouraged (Wreford et al., 2010). However, this is not easy as mitigation and adaptation may occur simultaneously, but differ in their spatial, timing and geographical characteristics (Smith and Olesen, 2010).

Amongst the number of policies affecting Climate Change mitigation and adaptation in grassland-based systems in Europe, the newly reformed EU Common Agricultural Policy (CAP), in principle, has made a decisive move towards promoting a greener and climatically friendlier EU agricultural sector. The new CAP has introduced direct payments associated to different practices that, in some cases, are expected to enhance GHG mitigation and adaptation to Climate Change. Namely, new payments within Pillar I associated to the diversification of crop rotations, maintaining permanent pastures and ensuring Ecological Focus Areas should be targeting, in part, climate-friendly or climate-smart agriculture. Permanent pasture maintenance is an important way to prevent N emissions through avoiding plough management and conversion of permanent grasslands into arable lands (EU Regulation 1307/2013). Leguminous species are mentioned explicitly in the areas of ecological interest (N-fixing species) but there is no special plan for their promotion. Other practices, such as those mentioned in previous sections, grazing, for example, is encouraged directly through the support of agroforestry systems and forests with fire risk areas (through the Rural Development Programme (Pillar II)), avoiding huge amounts of C release and through cross-compliance via for example promotion of good standards for animal welfare. Floristic biodiversity should also be encouraged but are not explicitly mentioned within the new PAC to safeguard ecosystem functioning against adverse climatic fluctuations.

The replacement of permanent grasslands by forage maize is no longer allowed by the CAP as penalties are included in the last CAP if destruction above 5% is present. The new CAP, however, does not explicitly address the worrying import of feed in grassland-based intensive systems. In fact, in some countries, this is still indirectly encouraged through additional payments to more intensive systems. The CAP has been blamed for distortion of global markets in this sense. Khatun (2012) points at the absence of tariffs for animal feed as a key driver for fueling EU cheap imports of animal feed from Latin America and consequently, for the effect on land use, land use change, and forestry (LULUCF) outside of the EU and, thereby preventing from a huge potential for mitigating climate change by reducing emissions from deforestation and forest degradation (REDD+ programme) outside Europe. Policies, hence, can create both positive and perverse incentives for mitigation or adaptation (Wreford et al., 2010). A number of recent studies (e.g. Lassaletta et al., in press in Spain) suggest that many European Countries, either assisted by specific regulations (e.g. Kyoto, CAP) or fuelled by market pressures, are displacing large amounts of GHG emissions from their national primary sectors (e.g. grassland-based farming) to other countries via agricultural goods importing. For example, cattle farming in Europe, whose feed system was traditionally based mainly on-farm forage (e.g. grass) production, in the recent decades has shift to heavily depend on cheap imported protein (e.g. soybean) from South America, resulting in a reduction of GHG emissions in the European GHG inventories but more than offsetting this potential mitigation by a consequential increase of GHG emissions by mainly land use change in South America. Much of these emissions are produced in non- Annex B countries and consequently, C leakage is being produced in Europe. Displacing agricultural productivity may indeed be an adaptation choice for countries, but this is certainly against securing Food Sovereignty and therefore, this jeopardizes the future resilience of the European food system. For example, if the conversion of annual crops to pasture is accompanied by a demand to grow annual crops

outside Europe, this would not represent a net mitigation but merely a shift in emissions and, in some cases, this would be an example of C and / or N₂O leakage.

Furthermore, a large part of the mitigation potential of grasslands is also subject to challenges in relation to effectiveness over different time-scales. For example, whereas certain types of mitigation activities (e.g. N₂O reduction from reduced N fertilization, CH₄ reduction in the rumen through animal diet changes, bioenergy) are effectively permanent since the emissions, once avoided, cannot be re-emitted (IPCC, 2014b), some activities that helped to sequester C (e.g. reducing tillage), can be reversible and non-permanent. Moreover, some of these practices to sequester soil C may also be constrained due to the saturation of grassland soils to sequester C indefinitely. Therefore protecting the large C stocks in grasslands should be an important management and policy target, rather than necessarily trying to increase the C stocks (Smith, in press) since it is easier and faster for soils to lose C than it is for them to gain C (Johnson et al. 2009).

Mitigation options for any of the GHG gases must also be tailored to the specific soil, climatic and production system conditions (Bustamante et al., in press). There will be very few strategies that are universally applicable for all systems and under any climatic circumstances. All mitigation options certainly affect and are affected by the cycles of C and N. Nitrogen and C cycles are also currently decoupled for most intensive grassland systems (Soussana and Lemaire, 2014), these systems release by ruminants bound-C digestible as CO₂ and CH₄, and return digestible N in high concentrations (urine patches). The coupling / decoupling of C and N makes an added difficulty to analyze the effectiveness of mitigating measures as sometimes some of the measures that increase soil C storage, for example, addition of manure, can also increase losses of N₂O by increasing soluble C in the system. In contrast, measures that promote the reduction of N₂O can cause a net loss of C from the system through increased soil respiration (Scholefield et al., 2005). Moreover, some of the mitigation methods lead to pollution swapping (e.g. NH₃ volatilization, leaching of NO₃⁻), and losses in biological diversity and / or productivity (Del Prado and Scholefield, 2008), and also can cause numerous interactions between mitigation measures so that their effect in the case of using multiple measures simultaneously are not necessarily additive (Del Prado et al., 2010).

Also, the reference unit to which GHG emissions relate within the CAP is commonly the forage area, which may not, in some cases, coincide with the preferred reference unit used by the agroindustry (C footprint or GHG per unit of product). The emphasis therefore seems to have been diverted from what the consumers and markets dynamics are essentially promoting. Preferably, one should consider more than one reference unit or functional unit (e.g. per hectare and per unit of output) at the same time to avoid conflicts of interpretation about what is true / false mitigation (Del Prado et al., 2010). Agroindustry generally uses the Life Cycle Assessment (LCA) as the methodology choice in order to report GHG emissions from the full cycle of the production of a food. A key element still unsolved is the way LCA assigns different amounts of GHG emissions to different goods according to its market-based value. Given a specific policy context, the farmer may choose among the most cost-effective and easier-to-adopt options. Ecosystem services which currently have no market value may become valuable also in monetary terms in the future. Some farmers may, therefore, in the future also seek to maximize the ecosystem service value. Alternative methodologies are already suggesting that, for products that are produced through extensive and in some cases greener conditions, these emissions should be split according to not only market but non-market (e.g. ecosystem services) values (Ripoll-Bosch et al., 2013) as well.

An important issue that may not be reflected in the new CAP and in other policies is the alarming growth tendency of feeding ruminants (e.g. dairy cattle: Del Prado et al., 2013b) with a greater amount of feed ingredients which could be used directly in the human food

chain (e.g. cereals) (Eisler et al., 2014). This relates very significantly to the potential competitive advantage that pasture-based livestock (ruminants generally are able to use low-quality plant biomass and that is inedible to humans) might have over another livestock (e.g. monogastric animals). Policies therefore should be useful to overturn this trend.

Additionally, non-climate policies and regulations are already in place for other environmental issues (e.g. water quality, NH₃) and have consistently assisted in reducing GHG emissions from the agricultural sector (e.g. EU: Velthof et al., 2014). Nitrate leaching losses, however, are expected to increase for numerous areas that are already constrained in their nutrient use by the EU Nitrates Directive (Anon, 1991) in Europe and for feed commonly used in animal diets, for example wheat (Olesen et al., 2007). This increase in NO₃- leaching may trigger more stringent regulations and hence affect animal productivity and GHG emissions, which may challenge climate change adaptation also from a policy perspective. Research-oriented policies should and already have a role, for example, in encouraging the study of new grass varieties that can better adapt to climate change and also present properties that can increase the efficiency of use of nutrients and energy in the soil-plant-animal system.

It is therefore imperative that all the policies, from the local to the global levels, are appropriately integrated with the policies relating to climate change, bioenergy, food, waste, research and health in order to promote a net reduction of GHG from the standpoint not only of production (supply) but also of demand in order to avoid possible market distortions and maladaptation practices at all levels.

Acknowledgements

This paper is a contribution to the FACCE MACSUR knowledge hub. MACSUR is funded through national funding bodies as part of the EU FACCE-JPI (Joint Programming Initiative for Agriculture, Climate Change, and Food Security). AGFORWARD and CICYT (Spanish Ministry) projects are also acknowledged.

6. References

- Alford A. R., Hegarty R. S., Parnell P. F., Cacho O. J., Herd R. M. and Griffith, G. R. (2006) The impact of breeding to reduce residual feed intake on enteric methane emissions from the Australian beef industry. *Australian Journal of Experimental Agriculture* 46, 813-820.
- Álvaro-Fuentes J., Easter M., Cantero-Martínez C. and Paustian K. (2011). Modelling soil organic carbon stocks and their changes in the northeast of Spain. *European Journal of Soil Science* 62(5), 685-695.
- Anon. (1991) Council Directive concerning the protection of water against pollution caused by nitrates from agricultural sources. *Official Journal of the European Communities* (91/676/EEC) 8572/91.
- Bates B., Kundzewicz Z. W., Wu S. and Palutikof J. (2008) Climate change and water. Intergovernmental Panel on Climate Change (IPCC).
- Bell M. J., Eckard R. J. and Cullen B. R. (2012) The effect of future climate scenarios on the balance between productivity and greenhouse gas emissions from sheep grazing systems. *Livestock Science* 147, 126-138.
- Betts R. A., Boucher O., Collins M., Cox P. M., Falloon P.D., Gedney N., Hemming D. L., Huntingford C., Jones C. D., Sexton D. M. H. and Webb M.J. (2007) Projected increase in continental runoff due to plant responses to increasing carbon dioxide. *Nature* 448, 1037-1041.
- Bindi M. and Olesen J. E. (2011) The responses of agriculture in Europe to climate change. *Regional Environmental Change* 11(1), 151-158.
- Brown L., Scholefield D., Jewkes E.C., Lockyer D.R. and Del Prado A. (2005) NGAUGE: A decision support system to optimize N fertilization of British grassland for economic and environmental goals. *Agriculture Ecosystems and Environment* 109, 20-39.
- Bryan E., Deressa T. T., Gbetibouo G. A. and Ringler C. (2009) Adaptation to climate change in Ethiopia and South Africa: options and constraints. *Environmental Science and Policy* 12(4), 413-426.
- Bryan B. A., King D. and Ward J. R. (2011) Modelling and mapping agricultural opportunity costs to guide landscape planning for natural resource management. *Ecological Indicators* 11, 199-208.
- Bustamante M., Robledo-Abad C., Harper R., Mbow C., Ravindranat N. H., Sperling F., Haberl H., De Siqueira Pinto A. and Smith P. (in press) Co-benefits, trade-offs, barriers and policies for greenhouse gas mitigation in the agriculture, forestry and other land use (AFOLU) sector. *Global Change Biology*.
- Cardenas L. M., Thorman R., Ashlee N., Butler M., Chadwick D., Chambers B., Cuttle S., Donovan N., Kingston H., Lane S., Dhanoa M. S. and Scholefield D. (2010) Quantifying annual N₂O emission fluxes from grazed grassland under a range of inorganic fertiliser nitrogen inputs. *Agriculture Ecosystems and Environment* 136, 218-226.
- Christensen J. H., Hewitson B., Busuioc A., Chen A., Gao X., Held R., Jones R., Kolli R. K., Kwon W. K., Laprise R., Magana Rueda V., Mearns L., Menendez C. G., Räisänen J., Rinke A., Sarr A., Whetton P., Arritt R., Benestad R., Beniston M., Bromwich D., Caya D., Comiso J., de Elia R. and Dethloff K. (2007) Regional climate projections, *Climate Change, 2007: The Physical Science Basis. Contribution of Working group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, University Press, Cambridge, Chapter 11.
- Correal E., Erena M., Ríos S., Robledo A. and Vicente M. (2009) Agroforestry systems in Southeastern Spain. In: Rigueiro-Rodríguez A., McAdam J. and Mosquera-Losada M.R. (eds) *Agroforestry in Europe*. Springer: 183-210.
- De Klein C. M. and Ledgard S. (2005). Nitrous Oxide Emissions from New Zealand Agriculture - key Sources and Mitigation Strategies. *Nutrient Cycling in Agroecosystems* 72, 77-85.

Del Prado A., Chadwick D., Cardenas L., Misselbrook T., Scholefield D. and Merino P. (2010) Exploring systems responses to mitigation of GHG in UK dairy farms. *Agriculture, Ecosystems and Environment* 136 (3-4): 318-332.

Del Prado A., Crosson P., Olesen J.E. and Rotz A. (2013a) Whole-farm models to quantify greenhouse gas emissions and their potential use for linking climate change mitigation and adaptation in temperate grassland ruminant-based farming systems. *Animal* 7 (suppl.2), 373-385.

Del Prado A., Mas K., Pardo G. and Gallejones. P. (2013b) Modelling the interactions between C and N farm balances and GHG emissions from confinement dairy farms in northern Spain. *Science of The Total Environment* 465, 156-165.

Del Prado A., Merino P., Estavillo J. M., Pinto M. and González-Murua, C. (2006) N₂O and NO emissions from different N sources and under a range of soil water contents. *Nutrient cycling in agroecosystems* 74(3), 229-243.

Del Prado A., Misselbrook T., Chadwick D., Hopkins A., Dewhurst R.J., Davison P., Butler A., Schröder J. and Scholefield D. (2011) SIMSDAIRY: A modelling framework to identify sustainable dairy farms in the UK. Framework description and test for organic systems and N fertiliser optimisation. *Science of the Total Environment* 409: 3993-4009.

Del Prado A. and Scholefield D. (2008). Use of SIMSDAIRY modelling framework system to compare the scope on the sustainability of a dairy farm of animal and plant genetic-based improvements with management-based changes. *Journal of Agricultural Science* 146: 1-17.

Del Prado A., Shepherd A., Wu L., Topp C., Moran D., Tolcamp B. and Chadwick D. (2009) Modelling the effect of climate change on environmental pollution losses from dairy systems in the UK. BC3 Working Paper Series 2009-07. Basque Centre for Climate Change (BC3). Bilbao, Spain

Dresner S., Ekins P., McGeevor K. and Tomei J. (2007) Forest and climate change global understandings and possible responses. In: Freer-Smith P.H., Boradmeadow M.S.J. and Lynch JM (eds.) *Forestry and climate change* CAB International, Wallingford, UK, pp 38-48

Dobbie K. E and Smith K. A. (2001) The effects of temperature, water-filled pore space and land use on N₂O emissions from an imperfectly drained gleysol. *European Journal of Soil Science* 52(4), 667-673.

Eagle A. J., Olander L. P. and Donald L. S. (2012) Chapter Three - Greenhouse Gas Mitigation with Agricultural Land Management Activities in the United States-A Side-by-Side Comparison of Biophysical Potential. *Advances in Agronomy*. Academic Press.

Eisler M. C., Lee M. R., Tarlton J. F., Martin G. B., Beddington J., Dungait J.A., Greathead H., Liu J., Mathew S., Miller H., Misselbrook T., Murray P., Vinod V. K., Van Saun R. and Winter M. (2014) Agriculture: Steps to sustainable livestock. *Nature* 507 (7490), 32.

Estavillo J. M., Merino P., Pinto M., Yamulki S., Gebauer G., Sapek A. and Corre W. (2002) Short term effect of ploughing a permanent pasture on N₂O production from nitrification and denitrification. *Plant and Soil* 239(2), 253-265.

European Union (EU) (2013) Communication from the Commission on an EU Forest Action Plan COM 2013 http://ec.europa.eu/agriculture/forest/strategy/communication_en.pdf

Graux A. I., Bellocchi G., Lardy R. and Soussana J. F. (2013) Ensemble modelling of climate change risks and opportunities for managed grasslands in France. *Agricultural and Forest Meteorology*, 170, 114-131.

Hart K. J., Yañez-Ruiz D. R., Duval S. M., McEwan N. R. and Newbold C. J. (2008) Plant extracts to manipulate rumen fermentation. *Animal Feed Science Technology* 147, 8-35.

Havlík P., Valin H., Herrero M., Obersteiner M., Schmid E., Rufino M. C., Mosnier A., Thornton P. K., Battcher H., Conant R. T., Frank S., Fritz S., Fuss S., Kraxner F. and Notenbaert A. (2014) Climate change mitigation through livestock system transitions. *Proceedings of the National Academy of Sciences* 111, 3709-3714.

Howden S. M., Crimp S. J., and Stokes C. J. (2008) Climate change and Australian livestock systems: impacts, research and policy issues. *Animal Production Science* 48(7), 780-788.

Johnson A. E., Poulton P.R and Coleman K. (2009) Soil organic matter: Its importance in sustainable agriculture and carbon dioxide fluxes. *Advances in Agronomy* 101, 1-57.

Intergovernmental Panel on Climate Change (IPCC) (2007) Climate change: the physical science basis. In: Solomon S., Qin D., Manning M., Chen Z., Marquis M., Averyt K.B., Tignor M. and Miller H.L. (eds.), Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change Chapter 2. Cambridge University Press, United Kingdom. 2007.

Intergovernmental Panel on Climate Change (IPCC) (2014a) Climate Change 2014: Impacts, Adaptation, and Vulnerability. IPCC WGII AR5 Food security and Food production systems. http://ipcc-wg2.gov/AR5/images/uploads/WGIIAR5-Chap7_FGDall.pdf.

Intergovernmental Panel on Climate Change (IPCC) (2014b) Climate Change 2014: Mitigation of Climate Change. IPCC WGIII AR5. Agriculture, Forestry and Other Land Use (AFOLU). http://report.mitigation2014.org/drafts/final-draft-postplenary/ipcc_wg3_ar5_final-draft_postplenary_chapter11.pdf

Khatun K. (2012) Reform or Reversal; Implications of the Common Agricultural Policy (CAP) on Land Use, Land Use Change and Forestry (LULUCF) in Developing Countries. *Conservation Letters* 5. 99-106.

Kreyling J., Thiel D., Simmnacher K., Willner E., Jentsch A. and Beierkuhnlein C. (2012) Geographic origin and past climatic experience influence the response to late spring frost in four common grass species in central Europe. *Ecography* 35(3), 268-275.

Kumar A. B. M. and Nair P. K. R. (2011) Carbon Sequestration Potential of Agroforestry Systems: Opportunities and Challenges. Springer.

Lal R. (2005) Forest soils and C sequestration. *Forest Ecology and Management* 220:242-258

Lal R., Delgado J. A., Groffman P. M., Millar N., Dell C. and Rotz A. (2011) Management to mitigate and adapt to climate change. *Journal of Soil and Water Conservation* 66(4), 276-285.

Lassaletta L., Aguilera E., Sanz-Cobena A., Pardo G., Billen G., Garnier J. and Grizzetti, B. (in press) Leakage of nitrous oxide emissions within the Spanish agro-food system in 1961-2009. *Mitigation and Adaptation Strategies for Global Change* 1-20.

Ledgard S.F., Menneer J.C., Dexter M.M., Kear M.J., Lindsey S., Peters J.S. and Pacheco D. (2008) A novel concept to reduce nitrogen losses from grazed pastures by administering soil nitrogen process inhibitors to ruminant animals: A study with sheep. *Agriculture, Ecosystems and Environment* 125, 148-158.

Liu C., Lu M., Cui J., Li B. and Fang C. (2014) Effects of straw carbon input on carbon dynamics in agricultural soils: a meta-analysis. *Global Change Biology* 20: 1366-1381.

Macleod C. J. A., Humphreys M. W., Whalley W. R., Turner L., Binley A., Watts C. W., Skot L., Joynes A., Hawkins S., King I. P., O'Donovan S. and Haygarth, P. M. (2013) A novel grass hybrid to reduce flood generation in temperate regions. *Scientific reports*, 3.

Misselbrook T., Del Prado A. and Chadwick D. (2013) Opportunities for reducing environmental emissions from forage-based dairy farms. *Agriculture and Food Science* 22: 93-107.

Monteny G. J., Bannink A. and Chadwick D. (2006) Greenhouse gas abatement strategies for animal husbandry. *Agriculture, Ecosystems and Environment* 112(2), 163-170.

Moreno G. and Pulido F. (2009) The functioning, management and persistence of Dehesas. In: Rigueiro-Rodríguez A., McAdam J. and Mosquera-Losada M.R. (eds) *Agroforestry in Europe*. Springer: 127-160.

Mosquera-Losada M. R., Freese D., Rigueiro-Rodríguez A. (2011a) Carbon Sequestration Potential of Agroforestry Systems: Opportunities and Challenges, *Advances in Agroforestry* 8, 43-59

Mosquera-Losada M. R., Morán-Zuloaga D. and Rigueiro-Rodríguez A. (2011b) Effects of lime and sewage sludge on soil, pasture production and tree growth in a six year old *Populus Canadensis* Moench silvopastoral system. *Journal of Plant Nutrition and Soil Science* 174:145-153.

Mosquera-Losada M. R., González-Rodríguez A. (1998) Effect of annual stocking rates in grass and maize+rye system on production by dairy cows. *Grass and Forage Science* 53 (2), 95-108.

Mountford J.O., Lakhani K.H. and Kirkham F.W. (1993) Experimental assessment of the effects of nitrogen addition under hay-cutting and aftermath grazing on the vegetation of meadows on a Somerset peat moor. *Journal of Applied Ecology* 30: 321-332.

Olesen J. E., Trnka M., Kersebaum K. C., Skjelvåg A. O., Seguin B., Peltonen-Sainio P., Rossi F., Kozyra J. and Micale F. (2011) Impacts and adaptation of European crop production systems to climate change. *European Journal of Agronomy* 34, 96-112.

Pan G., Smith P. and Pan W. (2009) The role of soil organic matter in maintaining the productivity and yield stability of cereals in China. *Agriculture, Ecosystems and Environment* 129(1), 344-348.

Papanastasis V. P., Yiakoulaki M. D., Decandia M. and Dini-Papanastasi O. (2008) Integrating woody species into livestock feeding in the Mediterranean areas of Europe. *Animal Feed Science and Technology* 140(1), 1-17.

Pinto M., Merino P., Del Prado A., Estavillo J. M., Yamulki S., Gebauer G. and Oenema O. (2004) Increased emissions of nitric oxide and nitrous oxide following tillage of a perennial pasture. *Nutrient Cycling in Agroecosystems* 70(1), 13-22.

Rinne M., Dragomir C., Kuoppala K., Marley C., Smith J. and Yañez-Ruiz D. (2012) Novel and underutilized feed resources - potential for use in organic and low input dairy production. In: Rahmann G. and Godinho D. (eds.) *Agriculture and Forestry Research* (362), pp. 404-407.

Rigueiro-Rodríguez A., McAdam J., Mosquera-Losada M. R. (2009) *Agroforestry in Europe*. Springer: 127-160.

Ripoll-Bosch R., de Boer I. J. M., Bernues A. and Vellinga T. V. (2013) Accounting for multi-functionality of sheep farming in the carbon footprint of lamb: A comparison of three contrasting Mediterranean systems. *Agricultural Systems* 116, 60-68.

Rosenzweig C. and Tubiello F. N. (2007) Adaptation and mitigation strategies in agriculture: an analysis of potential synergies. *Mitigation and Adaptation Strategies for Global Change* 12(5), 855-873.

Schmidhuber J. and Tubiello F. N. (2007) Global food security under climate change. *Proceedings of the National Academy of Sciences* 104(50), 19703-19708

Scholefield D., Jarvis S. C., Brown L., Del Prado A., Hopkins A. and Cardenas L. (2005) Feed back and Feed-Forward Interactions between Climate Change and Grassland-Based Agriculture. Avoiding Dangerous Climate Change. *Proceeding of the International Symposium on Stabilisation of Greenhouse Gases*, Met Office, Exeter, UK, 1-3 February 2005.

Smith P. (in press). Do grasslands act as a perpetual sink for carbon? *Global Change Biology*.

Smith P., Martino D., Cai Z., Gwary D., Janzen H., Kumar P, McCarl B., Ogle S., O'Mara F., Rice C., Scholes B., Sirotenko O., Howden M., McAllister T., Pan G., Romanenkov V., Schneider U. and Towprayoon (2007) Policy and technological constraints to implementation of greenhouse gas mitigation options in agriculture. *Agriculture, Ecosystems and Environment* 118(1), 6-28.

Smith P. and Olesen J. (2010) Synergies between the mitigation of, and adaptation to, climate change in agriculture. *The Journal of Agricultural Science* 148, 543-552.

Soussana, J.-F. & Lemaire, G. (2014) Coupling carbon and nitrogen cycles for environmentally sustainable intensification of grasslands and crop-livestock systems. *Agriculture, Ecosystems & Environment*, 190, 9-17.

Soussana J. F. and Lüscher A. (2007) Temperate grasslands and global atmospheric change: a review. *Grass and Forage Science* 62(2), 127-134.

Takács V. and Frank N. (2009) The traditions, resources and potential of forest growing and multipurpose shelterbelts in Hungary. In: Rigueiro-Rodríguez A, McAdam J, Mosquera-Losada M. R. (eds) *Agroforestry in Europe*. Springer: 415-438.

Theng B. K. G., Tate K. R., Sollins P., Moris N., Madkarni N. and Tate R. (1989) Constituents of organic matter in temperate and tropical soils. In: Coleman DC, Oades JM, Uehara G (eds.) *Dynamics of soil organic matter in tropical ecosystems*. College of Tropical agriculture and Human Resources, University of Hawaii, Honolulu, pp 5-32

Tingem M., Rivington M. and Bellocchi G. (2009) Adaptation assessments for crop production in response to climate change in Cameroon. *Agronomy for sustainable development* 29(2), 247-256.

UNEP (2013). Drawing down N₂O to protect Climate and the Ozone layer. A UNEP synthesis report. United Nations Environment Programme (UNEP), Nairobi, Kenya.
[http://www.unep.org/publications/ebooks/UNEPN₂O_report/](http://www.unep.org/publications/ebooks/UNEPN2O_report/)

Van den Pol-van Dasselaar A. (2012). Qualitative overview of mitigation and adaptation options and their possible synergies & trade-offs. Deliverable 8.1 of EU project AnimalChange, 32 p.

Van den Pol-van Dasselaar A., Vellinga T. V., Johansen A. and Kennedy E. (2008) To graze or not to graze, that's the question. *Grassland Science in Europe* 13: 706-716.

Vellinga T. V., Van den Pol-van Dasselaar A. and Kuikman P. J. (2004) The impact of grassland ploughing on CO₂ and N₂O emissions in the Netherlands. *Nutrient Cycling in Agroecosystems* 70(1), 33-45.

Vellinga T. V. and Hoving I. E. (2011) Maize silage for dairy cows: mitigation of methane emissions can be offset by land use change. *Nutrient cycling in agroecosystems* 89(3), 413-426.

Velthof G. L., Lesschen J. P., Webb J., Pietrzak S., Miatkowski Z., Pinto M., Kros J. and Oenema O. (2014) The impact of the Nitrates Directive on nitrogen emissions from agriculture in the EU-27 during 2000-2008. *Science of The Total Environment* 468, 1225-1233.

Velthof G. L., Hoving I. E., Dolfin J., Smit A., Kuikman P. J. and Oenema O. (2010). Method and timing of grassland renovation affects herbage yield, nitrate leaching, and nitrous oxide emission in intensively managed grasslands. *Nutrient Cycling in Agroecosystems* 86, 401-412.

Verchot L., M. van Noordwijk S., Kandji T., Tomich C., Ong A., Albrecht J., Mackensen C., Bantilan K., Anupama and Palm C. (2007) Climate change: linking adaptation and mitigation through agroforestry. *Mitigation and Adaptation Strategies for Global Change* 12:1381-1386.

Verge X. P. C., De Kimpe C. and Desjardins R. L. (2007) Agricultural production, greenhouse gas emissions and mitigation potential. *Agricultural and Forest Meteorology* 142(2), 255-269.

Volaire F., Barkaoui K. and Norton M. (2014) Designing resilient and sustainable grasslands for a drier future: Adaptive strategies, functional traits and biotic interactions. *European Journal of Agronomy* 52, 81-89.

Waghorn G. C. and McNabb W. C. (2003) Consequences of plant phenolic compounds for productivity and health of ruminants. *Proceedings of the Nutrition Society* 62, 383-392.

Wilkins P.W., Allen D.K. and Mytton L. R. (2000) Differences in the nitrogen use efficiency of perennial ryegrass varieties under simulated rotational grazing and their effects on nitrogen recovery and herbage nitrogen content. *Grass and Forage Science* 55, 69-76.

Wrage N., Velthof G. L., Van Beusichem M. L. and Oenema O. (2001) Role of nitrifier denitrification in the production of nitrous oxide. *Soil Biology and Biochemistry* 33(12), 1723-1732.

Wreford A., Moran D. and Adger N. (2010) *Climate Change and Agriculture: Impacts, Adaptation and Mitigation*. OECD Publishing, Paris, France.

Zaman M., Saggar S., Blennerhassett J. D. and Singh J. (2009) Effect of urease and nitrification inhibitors on N transformation, gaseous emissions of ammonia and nitrous oxide, pasture yield and N uptake in grazed pasture system. *Soil Biology and Biochemistry* 41, 1270-1280.

Zhang W. F., Dou Z., He P., Ju X.T., Powlson D., Chadwick D., Norse D., Lu Y. L., Zhang Y., Wu L., Chen X. P., Cassman K.G. and Zhang F.S. (2013). New technologies reduce greenhouse gas emissions from nitrogenous fertilizer in China. *PNAS* 110, 8375-8380.